

CollaMex is a 100% natural,
multifunctional bone, joint and
personal care ingredient

CollaMex

COLLAGEN CHONDROITIN COMPLEX



**WAITAKI
BIOSCIENCES**
A DIVISION OF PHARMAZEN LIMITED

CollaMex

is a 100%
natural source of
chondroitin sulphate
and type II collagen.

Multi-functional Ingredient

CollaMex is unique and proprietary natural ingredient for use in a wide range of formulations related to joint health, bone health and personal care products.

CollaMex contains naturally occurring **chondroitin sulphate** along with high concentrations of **type II collagen** peptides, bioactive **growth factors** and a range of **trace minerals**, all within a single ingredient package to deliver unsurpassed functionality.

Low Molecular Weight Chondroitin Sulphate

One of the group of compounds known as Glycosaminoglycan (GAG), chondroitin sulphate is present in the joint cartilage as part of the larger proteoglycan molecule, aggrecan.

Chondroitin sulphate provides the building blocks to support repair of damaged cartilage and interferes with the progression of structural changes in the joint tissue through several mechanisms. These include increased synthesis of joint proteoglycans^{1,2}, inhibition of matrix metalloproteases that can significantly degrade joint cartilage³ and anti-inflammatory properties. The overall result of these bioactivities is a reduction in the progression of joint structural changes and reduced pain⁴.

While chondroitin sulphate is better known for its use in joint health, research has also shown it has benefits in assisting with the regeneration of injured bone. Work by Brandt et al⁵ has demonstrated that the addition of chondroitin sulphate to bone cements significantly improved bone building properties.

Chondroitin sulphate is also known to be an effective ingredient in skin care formulations due to its ability to attract and retain water⁶.

CollaMex contains chondroitin sulphate in the range of 16,000 – 17,000 Daltons.

Studies have shown that the form (i.e. bovine versus shark) and molecular weight of chondroitin sulphate directly influences its absorption after oral administration. Lower molecular weight chondroitin sulphate of bovine origin has a higher gastrointestinal absorbability⁷.

The molecular weight range of the chondroitin sulphate in CollaMex has been shown to be within the range expected for bovine origin material with a peak at the lower molecular weight range of 16,000 - 17,000 Daltons.⁸

Type II Collagen

Type II collagen is the major structural component present in the extra cellular matrix (ECM) of connective tissues. Proper functioning of joint cartilage is dependent on the maintenance of the ECM, a process which is controlled by cartilage cells, known as chondrocytes.

Oral supplementation of type II collagen peptides (hydrolysed collagen) has been shown to help the repair of damaged cartilage by stimulating a dose dependent increase in type II collagen secretion by chondrocytes and increasing biosynthesis of proteoglycans⁹.

Proprietary, enzymatically controlled collagen hydrolysis

The specialised enzymatic manufacturing process carefully developed for CollaMex results in a finished powder with a high concentration of type II collagen peptides.

Specialised analysis of CollaMex shows a molecular weight range that includes collagen peptides of <8,000 Daltons.

Cartilage Derived Growth Factors

Waitaki has undertaken an ambitious project to identify growth factors present in CollaMex.

A range of growth factors were found to be present, in particular Insulin Like Growth Factor I (IGF I), but also Transforming Growth Factor β I (TGF β I) and Cartilage Derived Growth Factor (CDGF).

Gentle manufacturing process allows greater retention of the bioactives in CollaMex.

The Growth factors are a diverse group of proteins with a myriad of biological functions within the body. Because they are proteins they are very sensitive to harsh manufacturing processes, however the carefully controlled processing techniques used in the manufacture of CollaMex help to preserve these delicate entities within the finished powder.

IGF-I may have a dual benefit on compromised cartilage. Work by Neidel et al (1994)¹⁰ demonstrated that IGF-I partially prevented the inhibition of cartilage proteoglycan synthesis caused by Interleukin I (IL-1). In addition to this beneficial effect on the extracellular matrix, the growth factor actually accelerated recovery of proteoglycan production, after the free IL-1 was withdrawn.

CollaMex

Purity and functionality guaranteed.

CollaMex Features

- Gentle enzymatic manufacturing process preserves maximum biological activity
- Low molecular weight chondroitin sulphate for maximum absorbability
- High concentration of type II collagen peptides
- Independently tested source of growth factors to stimulate cartilage recovery
- Two convenient variants for maximum formulation flexibility
- CollaMex is manufactured from 100% safe and natural raw materials
- CollaMex contains no artificial additives or modifiers

High Quality Raw Materials

CollaMex is not just highly processed gelatin

Most collagen containing products are manufactured from inedible hides, and unspecified bone fragments that are left over from meat processing activities. Not surprisingly, these crude raw materials are subjected to harsh chemical treatments to sterilise and purify them to human consumption standards. These types of products are also known as gelatin and contain predominantly type I collagen. Other collagen products may be manufactured from caged, mass reared, chicken sternums.

CollaMex, however, is produced from New Zealand and Australian free range, grass fed, export quality bovine cartilage, that is exclusively collected to Waitaki's own strict specifications. Bovine cartilage supplies type II collagen - the same type of collagen that is found in the joints.

Furthermore, the raw materials used in the production of CollaMex are already certified as human food grade even before manufacturing begins. This means we do not need to use harsh solvents, acids or sterilisation procedures during manufacture, and ensures maximum bioactivity is retained in the finished product.

1 Brand, 2 Ingredients

To offer the highest degrees of flexibility for formulators, Waitaki offers CollaMex in two forms:

CollaMex Original - a minimally processed powder that maintains maximum levels of bioactivity

- < 250 micron, fine free flowing powder
- Minimum 20% Chondroitin Sulphate
- Minimum 40% type II collagen
- Naturally occurring growth factors
- Ideally suited to encapsulation and tableting and topical personal care products (e.g. lip balm)

CollaMex Soluble - a 100% water soluble fine grade powder for greater formulation flexibility

- < 150 micron, fine free flowing powder
- 100% soluble in water
- 20% - 30% chondroitin sulphate
- Minimum 40% type II collagen
- Naturally occurring growth factors
- Ideally suited to dry mix beverage applications, tableting and encapsulation

References:

¹ Bassler C., Henrotin Y, and Franchimont P, In vitro evaluation of drugs proposed as chondroprotective agents. Int j Tissue React 1992; 14: 231-241.

² Glade MJ., Polysulfated glycosaminoglycan accelerates net synthesis of collagen and glycosaminoglycan by arthritic equine cartilage tissues and chondrocytes. American Journal of Veterinarian Research 1990; 51(5): 779-785.

³ Bartolucci C., et al, Inhibition of human leucocyte elastase by chemically and naturally oversulphated galactosaminoglycans. Carbohydrate Research, 1995;276:401-408.

⁴ Montfort J., et al, Biochemical basis of the effect of chondroitin sulphate on osteoarthritis articular tissues. Ann.rheum. Dis.2008;67:735-740.

⁵ Brandt JH., et al, Biometric mineralization of collagen by combined fibril assembly and calcium phosphate formation. Chem. Matter. 1999;11:2694-2701.

⁶ Eskelinin A., Santalahati J., Special natural cartilage polysaccharide for the treatment fo sun damaged skin in females. Journal International Medical Research 1992; 20(2):99-105.

⁷ Neil KM., et al., The role of glucosamine and chondroitin sulphate in treatment for and prevention of osteoarthritis in animals. Journal of the American Veterinary Association, 2005. 226;7:1079-1088.

⁸ Scott D., Slim G., Molecular Sieve Chromatography of Waitaki Biosciences Soluble Bovine Tracheal Cartilage. Industrial Research Ltd Report. April 2001.

⁹ Bello AE., Osser S., Collagen hydrolysate for the treatment of osteoarthritis and other joint disorders: a review of the literature. Current Medical Research and Opinion. 2006;22(11):2221-2231.

¹⁰ Neidel M., Schulze M., Soya insulin like growth factor accelerates recovery of articular cartilage proteoglycan synthesis in culture after inhibition by interleukin 1. Arch. Orthop. Trauma Surg. 1994;114:3-48